

METEOROID STRUCTURE: CURRENT VIEWS, ASTROPHYSICAL IMPORTANCE AND SOFIA POSSIBILITIES

Robert L. Hawkes¹

¹*Physics Department, Mount Allison University, Sackville, NB Canada E4L 1E6*

ABSTRACT

While recent meteor electro-optical observations and theoretical ablation models have provided clues to the structure of meteoroids and the role of volatile, possibly organic, materials in meteors, many questions remained unanswered. A SOFIA airborne platform offers promise for resolution of some of the key questions about meteoroid structure, and ultimately will contribute to questions about astrophysical origins and space operations hazards from natural meteoroids. The extension of the optical window for meteor studies into the near and mid-infrared regions, the relative freedom from atmospheric scintillation based noise, and the ability to observe meteors at low elevation angles are the key advantages to observations from the SOFIA upper deck platform.

INTRODUCTION

After decades of intensive work by numerous researchers, a definitive view of the structure and composition of meteoroids, and the mode of atmospheric ablation of these objects, remains elusive. There is general support for the idea that many cometary meteoroids have some sort of dustball structure in which a meteoroid is actually a collection of at least hundreds to thousands of small constituent grains (Jacchia 1955; Verniani 1969; Hawkes and Jones 1975; Fisher et al. 2000). Hawkes and Jones (1975) proposed a quantitative dustball model in which these constituent grains were bound by a more volatile second component. This proposed meteoroid structure is, in broad features at least, consistent with models of organic coated silicate interstellar and cometary grains proposed by Greenberg and others, and also consistent with the Halley dust observations of significant CHON composition. According to this model what we call a single meteor event may be a cluster of fundamental grains which have been released prior to intensive ablation (following loss of the volatile component high in the atmosphere), or one experience simultaneous grain release and intensive grain. Several authors have shown that the quantitative dustball model matches meteor observations (see e.g. Beech 1984). More recently Borovicka et al. (1999) have proposed a modification of the model in which Na plays a key role as the more volatile constituent.

If a sufficiently detailed picture of the composition and physical structure of these dustball meteoroids can be confirmed (e.g. the mass distribution of the constituent grains; the fraction of the original material was volatile; the degree of variability from meteoroid stream to meteoroid stream), then this will help to constrain models of solar system formation. The importance of delivery of organic material to the primitive and current Earth is also dependent on these issues (e.g. Jenniskens et al. 2000). Understanding meteor fragmentation and ablation processes is key to dealing with selection biases inherent in radar meteor detection (e.g. Campbell and Jones 2003). Furthermore, a clear understanding of the physical structure and ablation mode is key to a complete understanding of the potential hazards to space operations posed by natural meteoroids.

METEOROID FRAGMENTATION OVERVIEW

Several models of potential meteoroid structure and fragmentation have been proposed. One possibility is a compact, high strength structure (similar to stony meteorites) which fragments into a relatively small number of pieces due to a combination of thermal and aerodynamic stresses. The Peekskill meteorite fireball (see Fig. 1) graphically demonstrates this mechanism (Brown et al. 1994) which is common for bright meteors but appears to be rare in the size range studied with image intensified video techniques.




Fig. 1. In-flight fragmentation of the Peekskill meteorite fireball (from Brown et al. 1994).

A second mode of fragmentation is spraying of droplets from a molten surface of the meteoroid during atmospheric entry. A third possibility is that the meteor will fragment into smaller and smaller pieces during atmospheric flight in response to steadily growing dynamic pressures. A fourth possibility is continuous fragmentation through dispersal of many small grains during the atmospheric flight, and Babadzhanov (2002) argues that this is the dominant mode of fragmentation at least in the size range studied using photographic techniques. A fifth possibility is that the meteoroid is already fragmented into a collection of constituent grains prior to atmospheric entry. Of course combinations of these models are possible, for example a dustball meteor which has one or more episodes of fragmentation into large number of grains (and subsequent flare production) as well as continual release of smaller numbers of grains. In the remaining portions of this paper we will overview some of the questions related to meteoroid structure and meteor ablation which are particularly suited to contributions from SOFIA.




Fig. 2. Detailed meteor light curve obtained by background subtraction and line profile analysis from a wide field image intensified CCD system. Most of the apparent fluctuations are not true light curve fluctuations by rather instrumental noise and sky background variability (from Hawkes et al. 2001).

ARE METEOR LIGHT CURVES SMOOTH?

Perhaps the most obvious way to search for evidence of a dustball structure and fragmentation is to look for short scale irregularities on meteor light curves resulting from changes in ablation rate and surface area due to in-flight fragmentation. Jiang and Hu (2001) claim several detections demonstrating this effect and consistent with the Hawkes and Jones (1975) dustball model of meteoroid structure. Hawkes et al. (2001) proposed several techniques for doing this and reported on tentative early results. For example, if the stellar and sky background is subtracted from the meteor image, and an intensity plot profile is then taken, results such as those shown in Fig. 2 are obtained. At first glance this seems to confirm dustball fragmentation through light curve irregularities. However, the

combination of readout noise in the CCD, scintillation due to the atmosphere, image intensifier noise, and short temporal scale skyglow variations make it very difficult to isolate true short term.

Using observations from SOFIA two of these effects (atmospheric scintillations and skyglow temporal variations) will be significantly reduced. Two low noise image intensified CCD detectors with moderate to high spatial resolution should be employed in coincidence mode for this study. Meteor flares can be used to estimate the mass distribution of simultaneously released dustball grains (see e.g. Hawkes et al. 2002 and references therein). Unfortunately these are rare events for any single detectors. As Jenniskens et al. (2004) point out, the meteor rate from an airborne platform can be significantly enhanced by looking at very low elevations not possible for terrestrial observations.

IS METEOR WAKE DUE TO DIFFERENTIAL DECELERATION COMMON?

Meteor wake can be caused by long duration photochemical processes and by differential deceleration of grains with different physical characteristics so that light is simultaneously produced from a distributed spatial region. It has been difficult to obtain many clear examples of this with three previous studies of wake in faint meteors each providing only a few examples (Robertson and Hawkes 1992; Shadbolt and Hawkes 1992; Fisher et al. 2000). To have success in this it will be necessary to improve spatial and temporal resolution, while also reducing noise. Again, the reductions in scintillation and the darkening of the unresolved sky background may make SOFIA a good platform for this, particularly if one uses digital gated image intensifier technology connected to high spatial resolution optics (Hawkes et al. 2004).




Fig. 3. Two meteor images obtained using a gated image intensifier fibre-optically coupled to a digital CCD camera. The system is gated with an on:off ratio of 1:4 in both cases. Fragmentation induced wake eliminates most of the dark segments in the left hand image, while wake is of only minor importance in the right hand image (from Hawkes et al. 2004 in preparation).

There are several reported cases of clearly separated but clustered meteors (e.g. Watanabe et al. 2003). The study of these rare events could be enhanced from SOFIA due to darker backgrounds, and the possibility of looking at low elevation angles for a larger collecting area. Reduced scintillations from the SOFIA platform may assist with detection of dual or multiple fragment simultaneous meteors.

WHAT IS THE NATURE OF THE ORGANIC VOLATILE COMPONENT OF METEOROIDS?

Perhaps the most fundamental question is the nature of the volatile component of the quantitative dustball structure (Hawkes and Jones 1975). SOFIA offers the opportunity to extend the spectral range that can be

observed, which may offer opportunities to clearly identify the products of the organic components of meteoroids. This topic is covered in detail by Trigo-Rodriguez (2004) and Jenniskens et al. (2000). Study of the early parts of meteor light curves, which should correspond to ablation of volatile possibly organic material (or sputtered material – see below) are particularly important. The dark sky background offered by the SOFIA platform would assist with detection of these faint, possibly infrared rich signatures.

Until recently it has been assumed that little ablation occurs until the meteor reaches intensive vaporization temperatures. Coulson and Wikckramasinghe (2003) and Rogers et al. (2004) have shown that in some cases sputtering should be considered. Direct experimental verification of sputtering will need to come from detection of a low luminosity component in the early part of meteor light curves. This is very challenging observationally, but the reduced skyglow and scintillations from SOFIA will assist.

DISCUSSION

SOFIA Upper Deck offers excellent opportunities for electro-optical meteor observations, and the potential to definitively answer key questions about the structure and atmospheric ablation of meteoroids. In turn these will allow us to answer questions about our origins, in particular the dust conditions in the primitive solar system and the importance of delivery mechanisms of organic material to Earth through meteoroid ablation. The features and frequency of clustered dustball meteoroids are important for evaluating the hazards to space operations from meteoroid impacts.

Several key advantages of the SOFIA Upper Deck platform over ground based meteor observations are briefly summarized below.

- The spectral region which can be observed is expanded, in particular into the infrared where organic volatile components may be detected.
- The Airborne Leonid missions have illustrated that the enhanced collection area offered by low altitude pointing directions, only possible from airborne and space detectors, lead to significantly higher meteor rates. This allows more efficient collection of rare observational data such as clustered meteoroids, flare events, and spectra from a wider sampling of cometary meteoroids.
- The reduced atmospheric scintillation (and to a lesser degree the reduced unresolved skyglow background) offer the opportunity to search for slight irregularities on light curves as indicators of fragmenting dustballs.

Acknowledgements: I would like to acknowledge helpful discussions on issues related to this paper with the following current and recent research students in our meteor laboratory at Mount Allison (David Babcock, Jason Bussey, Margaret Campbell, Ashley Faloon, Amy Fisher, Kyle Hill, Andrew LeBlanc, Stephanie MacPhee, Ian Murray, Laura Parker, Chris Pollock, Leslie Rogers, Lloyd Taggart, and John Thaler, as well as collaborators from the University of Western Ontario (including Peter Brown, Jim Jones and Nicole Kaiser). The research of which this forms a part is funded by grants from the Natural Sciences and Engineering Research Council of Canada (NSERC).

REFERENCES

- Babadzhanov, P.B. Fragmentation and densities of meteors. *Astron. Astrophys* **384**, 317-321, 2002.
- Beech, M. The structure of meteoroids. *Mon. Not. R. astr. Soc.* **211** 617-620, 1984.
- Beech Martin and Murray, I.S. Leonid meteor light-curve synthesis. *Mon. Not. R. astr. Soc* **345** 696-704, 2003.
- Borovicka, J., Stork, R. and Bocek, J. First results from spectroscopy of 1998 Leonid meteors. *Meteoritics Planetary Sci.* **34**, 987-994, 1999.
- Brown, P., Ceplecha, Z., Hawkes, R.L., Wetherill, G., Beech, M. and Mossman, K. The orbit and atmospheric trajectory of the Peekskill meteorite from video records. *Nature* **367** 624-626, 1994.
- Campbell-Brown, M. and Jones, J. Determining the initial radius of meteor trains: fragmentation. *Mon. Not. R. astr. Soc* **343** 775-780, 2003.

- Campbell, M.D., Brown, P.G., Leblanc, A.G., Hawkes, R.L., Jones, J., Worden, S.P. and Correll, R.R. Image-intensified video results from the 1998 Leonid shower: I. Atmospheric trajectories and physical structure. *Meteoritics and Planetary Science* **35** 1259-1267, 2000.
- Coulson, S.G. and Wickramasinghe, N.C. Frictional and radiation heating of micron-sized meteoroids in the Earth's upper atmosphere. *Mon. Not. R. astr. Soc.* **343**, 1123-1130, 2003.
- Fisher, A.A., Hawkes, R.L., Murray, I.S., Campbell, M.D. and LeBlanc, A.G. Are meteoroids really dustballs? *Planetary and Space Science* **48** 911-920, 2000.
- Fujiwara, Y., Ueda, M., Shiba, Y., Sugimoto, M., Kinoshita, M., Shimoda, C. and Nakamura, T. Meteor luminosity at 160 km altitude from TV observations for bright Leonid meteors. *Geophys. Res. Lett.* **25**, 285-288, 1998.
- Hawkes, R.L., Brown, P.G., Kaiser, N.R., Faloon, A.J., Hill, K.A., Rogers, L.A. High Spatial and Temporal Resolution Optical Search for Evidence of Meteoroid Fragmentation (in preparation for Meteoroids 2004 Conference)
- Hawkes, R.L., Bussey, J.E., Macphee, S.L., Pollock, C.S. and Taggart, L.W. Techniques for high resolution meteor light curve investigations. *Meteoroids 2001 in ESA SP-495*, 281 - 286 , 2001
- Hawkes, R.L., Campbell, M.D., LeBlanc, A.G., Parker, L., Brown, P., Jones, J., Worden, S.P., Correll, R.R., Woodworth, S.C., Fisher, A.A., Gural, P., Murray, I.S., Connors, M., Montague, T., Jewell, D. and Babcock, D.D. The Size of Meteoroid Constituent Grains: Implications for Interstellar Meteoroids. in *Dust in the Solar System and Other Planetary Systems (Pergamon: Amsterdam, The Netherlands)* (ed. S.F. Green, I. Williams, T. McDonnell & N. McBride) 23-26, 2002.
- Hawkes, R.L. and Jones, J. A quantitative model for the ablation of dustball meteors. *Mon. Not. R. astr. Soc* **173** 339-356, 1975.
- Jacchia, L.G. The physical theory of meteors VIII Fragmentation as cause of the faint-meteor anomaly. *Astrophys. J.* **121**, 521-527, 1955.
- Jenniskens, P., Jost, H., Russell, R.W., Taylor, M.J., Castellano, T., Stenbaek-Nielson, H.C., and Rietmeijer, F.J.M., Towards a SOFIA Upper Deck Research Facility. Proceedings SOFIA Upper Deck Science Opportunities Workshop, 2004
- Jenniskens P., Wilson, M.A., Packan D., Laux, C.O., Krüger, C.H., Boyd, I.D., Popova, O.P. and Fonda, M. Meteors: A Delivery Mechanism of Organic Matter to the Early Earth. *Earth Moon and Planets* **82** 57-70, 2000.
- Jiang, X. and Hu, J. High resolution Leonid meteor light curves: implied meteoroid structure. *Planetary and Space Science* **49** 1281-1283, 2001.
- Koten, P., Spurny, P., Borovicka, J. and Stork, R. Extreme beginning heights for non-Leonid meteors. in *Meteoroids 2001 (ed. B. Warmbein)* **ESA SP-495**, 119-122, 2001.
- LeBlanc, A.G., Murray, I.S., Hawkes, R.L., Worden, P., Campbell, M.D., Brown, P., Jenniskens, P., Correll, R.R., Montague, T. and Babcock, D.D. Evidence for transverse spread in Leonid meteors. *Mon. Not. R. astr. Soc* **313** L9-L13, 2000.
- Robertson, M.C. and Hawkes, R.L. Wake in faint television meteors. *Asteroids, Comets, Meteors 1991* 517-520, 1992
- Rogers, L.A., Hill, K.A. and Hawkes, R.L. Mass loss due to sputtering and thermal processes in meteor ablation. *Planetary Space Sci.*, , 2004 (submitted)
- Shadbolt, L. and Hawkes, R.L. Absence of Wake in Faint Television Meteors. *Earth Moon and Planets* **68** 493-502, 1995.
- Spurny P., Betlem H., van't Leven J. and Jenniskens P. Atmospheric behavior and extreme beginning heights of the 13 brightest photographic Leonids from the ground-based expedition to China. *Meteoritics and Planetary Science* **35** 243-249, 2000.
- Trigo-Rodriguez, T.M. Detection of volatiles in cometary meteoroids using IR meteor spectroscopy Proceedings SOFIA Upper Deck Science Opportunities Workshop, 2004.
- Verniani, F. Structure and fragmentation of meteoroids. *Space Sci. Rev.* **10**, 230-261, 1969.
- Watanage, J-I, Tabe, I., Hasegawa, H., Hashimoto, H., Furse, T., Yoshikawa, M., Abe, S., Suzuki, B., Meteoroid clusters in Leonids: Evidence for fragmentation in space. *Publ. Astron. Soc. Japan* **55** L23-L26, 2003.

E-mail address of R. Hawkes: rhawkes@mta.ca

Manuscript received 2004 June 15, revised 2004 August 5, accepted 2004 August 5.